Author index

Abidin, İ. *see* Yargiçoğlu, P.
Ağar, A. *see* Yargiçoğlu, P.
Aizman, O. *see* Andersson, R.M.
Akai, M. *see* Kawashima, N.
Al-Khalili, L., Krämer, D., Wretenberg, P. & Krook, A. Human skeletal muscle cell differentiation is associated with changes in myogenic markers and enhanced insulin-mediated MAPK and PKB phosphorylation, 395
Andersen, O.K. *see* Terkelsen, A.J.
Andersson, G. *see* Zhang, Q.
Andersson, K.B. *see* Florholmen, G.
Andersson, R.M., Aizman, O., Aperia, A. & Brismar, H. Modulation of Na^+,K^+-ATPase activity is of importance for RVD, 329
Aoki, K., Kondo, N., Shimomura, Y., Yokoi, M., Iwanaga, K., Harada, H. & Katsuura, T. Sweating responses during activation of the muscle metaboreflex in humans is altered by time of day, 63
Aperia, A. *see* Andersson, R.M.
Arnolda, L.F. *see* Holobotovskyy, V.V.
Assimakopoulos, S.F., Vagianos, C.E., Patsoukis, N., Georgiou, C., Nikolopoulou, V. & Scopa, C.D. Evidence for intestinal oxidative stress in obstructive jaundice-induced gut barrier dysfunction in rats, 177
Austbo, B. *see* Florholmen, G.

Bäckström, T. & Franco-Cereceda, A. Intravasal microdialysis is superior to intramyocardial microdialysis in detecting local ischaemia in experimental porcine myocardial infarction, 5
Bányász, T. *see* Fülöp, L.
Beltman, J.G.M., Sargeant, A.J., Haan, H., van Mechelen, W. & de Haan, A. Changes in PCr/Cr ratio in single characterized muscle fibre fragments after only a few maximal voluntary contractions in humans, 187
Berg, A. *see* Bronstad, A.
Bigard, A.X. *see* Bricout, V.-A.
Billeter, R. *see* Wittwer, M.
Bilmen, S. *see* Yargiçoğlu, P.
Boemke, W. *see* Francis, R.C.E.
Breier, B.H. *see* Holt, R.I.G.
Bricout, V.-A., Serrurier, B.D. & Bigard, A.X. Clenbuterol treatment affects myosin heavy chain isoforms and MyoD content similarly in intact and regenerated soleus muscles, 271
Brismar, H. *see* Andersson, R.M.
Bronstad, A., Reed, R.K. & Berg, A. Effect of the cytoskeletal fixation agent phalloidin on transcapillary albumin transport and interstitial fluid pressure following subdermal prostaglandin E_1 administration in the rat, 125

Chen, R., Penny, D.J., Greve, G. & Lab, M.J. Rate dependence of mechanically induced electrophysiological changes in right ventricle of anaesthetized lambs during pulmonary artery occlusion, 13
Christensen, G. *see* Florholmen, G.
Clausen, T., Overgaard, K. & Nielsen, O.B. Evidence that the Na^+-K^+ leak/pump ratio contributes to the difference in endurance between fast- and slow-twitch muscles, 209
Cohen, K.D. *see* Murrant, C.L.

De Blasio, B.F., Røttingen, J.-A., Sand, K.L., Giaever, I. & Iversen, J.-G. Global, synchronous oscillations in cytosolic calcium and adherence in bradykinin-stimulated Madin-Darby canine kidney cells, 335
Delgado, J. *see* Morán, M.
Desplanches, D., Ecochard, L., Sempore, B., Mayet-Sornay, M.-H. & Favier, R. Skeletal muscle HSP72 response to mechanical unloading: influence of endurance training, 387
Deussen, A. *see* Loncar, R.
Duhamel, T.A. *see* Schertzer, J.D.
Duza, T. *see* Murrant, C.L.

Ecochard, L., Roussel, D., Sempore, B. & Favier, R. Stimulation of HSP72 expression following ATP depletion and short-term exercise training in fast-twitch muscle, 71
Ecochard, L. *see* Desplanches, D.
Ekberg, A. *see* Tesch, P.A.
Ekblom, B. *see* Malm, C.
Ekblom, Ö. *see* Malm, C.

Fall, C.H.D. *see* Holt, R.I.G.
Favier, R. *see* Desplanches, D.
Favier, R. *see* Ecochard, L.
Fihn, B.-M. *see* Sun, Y.
Flesche, C.W. *see* Loncar, R.
Florholmen, G., Andersson, K.B., Yndestad, A., Austbo, B., Henriksen, U.L. & Christensen, G. Leukaemia inhibitory factor alters expression of genes involved in rat cardiomyocyte energy metabolism, 133
Flück, M. *see* Wittwer, M.
Fowles, J.R. *see* Schertzer, J.D.
Fraga, S. *see* Magro, F.
Francis, R.C.E., Höhne, C., Kaczmarczyk, G. & Boemke, W. Effect of angiotensin II and endothelin-1 receptor blockade on the haemodynamic and hormonal changes after acute blood loss and after retransfusion in conscious dogs, 319
Franco-Cereceda, A. *see* Bäckström, T.
Fülöp, L., Bányász, T., Magyar, J., Szentandrássy, N., Varró, A. & Nánási, P.P. Reopening of L-type calcium channels in human ventricular myocytes during applied epicardial action potentials, 39

Galbo, H. *see* Stallknecht, B.
Georgiou, C. *see* Assimakopoulos, S.F.
Giaever, I. *see* De Blasio, B.F.
Gluckman, P.D. *see* Holt, R.I.G.
González, B. *see* Morán, M.
Green, H.J. *see* Schertzer, J.D.
Greve, G. *see* Chen, R.
Gümüşlü, S. *see* Yargiçoğlu, P.

de Haan, A. *see* Beltman, J.G.M.
Haan, H. *see* Beltman, J.G.M.
Hafting, T. *see* Haug, T.M.
Hansen, J. *see* Terkelsen, A.J.
Hansen, O. Quantification of α-subunit isoforms of Na,K-ATPase in rat resistance vessels, 49
Harada, H. *see* Aoki, K.

© 2004 Scandinavian Physiological Society

Haug, T.M., Hafting, T. & Sand, O. Inhibition of BK channels contributes to the second phase of the response to TRH in clonal rat anterior pituitary cells, 347
Helge, J.W. *see* Stallknecht, B.
Henriksen, U.L. *see* Florholmen, G.
Henriksson, J. Editorial, 1
Hogan, D.E. *see* McHugh, M.P.
Höhne, C. *see* Francis, R.C.E.
Holobotovskyy, V.V., Arnolda, L.F. & McKitrick, D.J. Effect of anaesthetic and rat strain on heart rate responses to simulated haemorrhage, 29
Holt, R.I.G., Syddall, H.E., Phillips, D.I.W., Martyn, C.N., Gluckman, P.D., Breier, B.H., Wood, P.J. & Fall, C.H.D. Serum insulin-like growth factor-I concentrations in late middle age: no association with birthweight in three UK cohorts, 359
Hoppeler, H. *see* Wittwer, M.

Inashima, S. *see* Yamada, T.
Ishido, M., Kami, K. & Masuhara, M. Localization of MyoD, myogenin and cell cycle regulatory factors in hypertrophying rat skeletal muscles, 281
Iversen, J.-G. *see* De Blasio, B.F.
Iwanaga, K. *see* Aoki, K.

Jensen, T.S. *see* Terkelsen, A.J.
Jodal, M. *see* Sun, Y.

Kaczmarczyk, G. *see* Francis, R.C.E.
Kajihara, H. *see* Yamada, T.
Kami, K. *see* Ishido, M.
Kano, Y., Sampei, K. & Matsudo, H. Time course of capillary structure changes in rat skeletal muscle following strenuous eccentric exercise, 291
Karelov, A. *see* Segerdahl, M.
Katsuura, T. *see* Aoki, K.
Kawashima, N., Nakazawa, K., Yamamoto, S.-I., Nozaki, D., Akai, M. & Yano, H. Stretch reflex excitability of the anti-gravity ankle extensor muscle in elderly humans, 99
Kiens, B. *see* Stallknecht, B.
Kim, M.B. *see* Murrant, C.L.
Kondo, N. *see* Aoki, K.
Krämer, D. *see* Al-Khalili, L.
Krook, A. *see* Al-Khalili, L.
Kurata, Y. *see* Shibamoto, T.

Lab, M.J. *see* Chen, R.
Lindberg, L.G. *see* Zhang, Q.
Lindquist, D.M. *see* Tesch, P.A.
Ling, Y.-Q. *see* Shibamoto, T.
Loncar, R., Flesche, C.W. & Deussen, A. Myocardial ferritin content is closely related to the degree of ischaemia, 21

McHugh, M.P. & Hogan, D.E. Effect of knee flexion angle on active joint stiffness, 249
McKitrick, D.J. *see* Holobotovskyy, V.V.
Magro, F., Fraga, S., Ribeiro, T. & Soares-da-Silva, P. Decreased availability of intestinal dopamine in transmural colitis may relate to inhibitory effects of interferon-γ upon L-DOPA uptake, 379
Magyar, J. *see* Fülöp, L.
Malm, C., Ekblom, Ö. & Ekblom, B. Immune system alteration in response to two consecutive soccer games, 143
Manso, R. *see* Morán, M.
Martyn, C.N. *see* Holt, R.I.G.
Masuhara, M. *see* Ishido, M.
Matsudo, H. *see* Kano, Y.
Matsunaga, S. *see* Yamada, T.
Mayet-Sornay, M.-H. *see* Desplanches, D.
van Mechelen, W. *see* Beltman, J.G.M.
Megías, A. *see* Morán, M.
Mølgaard, H. *see* Terkelsen, A.J.
Morán, M., Delgado, J., González, B., Manso, R. & Megías, A. Responses of rat myocardial antioxidant defences and heat shock protein HSP72 induced by 12 and 24-week treadmill training, 157
Moyna, N.M. & Thompson, P.D. The effect of physical activity on endothelial function in man, 113
Murrant, C.L., Duza, T., Kim, M.B., Cohen, K.D. & Sarelius, I.H. Arteriolar dilations induced by contraction of hamster cremaster muscle are dependent on changes in endothelial cell calcium, 231

Nakazawa, K. *see* Kawashima, N.
Nánási, P.P. *see* Fülöp, L.
Nara, I. *see* Yamada, T.
Narushima, M. *see* Shibamoto, T.
Nielsen, O.B. *see* Clausen, T.
Nikolopoulou, V. *see* Assimakopoulos, S.F.
Nozaki, D. *see* Kawashima, N.

Ogura, T. *see* Shibamoto, T.
Overgaard, K. *see* Clausen, T.

Patsoukis, N. *see* Assimakopoulos, S.F.
Penny, D.J. *see* Chen, R.
Phillips, D.I.W. *see* Holt, R.I.G.

Ravesloot, J.H. *see* Verkerk, A.O.
Reed, R.K. *see* Brønstad, A.
Ribeiro, T. *see* Magro, F.
Richter, E.A. *see* Stallknecht, B.
Røttingen, J.-A. *see* De Blasio, B.F.
Roussel, D. *see* Ecochard, L.

Sampei, K. *see* Kano, Y.
Sand, K.L. *see* De Blasio, B.F.
Sand, O. *see* Haug, T.M.
Sarelius, I.H. *see* Murrant, C.L.
Sargeant, A.J. *see* Beltman, J.G.M.
Schertzer, J.D., Green, H.J., Fowles, J.R., Duhamel, T.A. & Tupling, A.R. Effects of prolonged exercise and recovery on sarcoplasmic reticulum Ca^{2+} cycling properties in rat muscle homogenates, 195
Scopa, C.D. *see* Assimakopoulos, S.F.
Segerdahl, M. & Karelov, A. Experimentally induced ischaemic pain in healthy humans is attenuated by the adenosine receptor antagonist theophylline, 301
Sempore, B. *see* Desplanches, D.
Sempore, B. *see* Ecochard, L.
Serrurier, B.D. *see* Bricout, V.-A.
Shibamoto, T., Narushima, M., Ling, Y.-Q., Shimo, T., Tsuchida, H., Kurata, Y. & Ogura, T. Different hepatic vascular response to noradrenaline and histamine between guinea-pig and rat, 255
Shimo, T. *see* Shibamoto, T.
Shimomura, Y. *see* Aoki, K.
Sjövall, H. *see* Sun, Y.
Soares-da-Silva, P. *see* Magro, F.
Stallknecht, B., Kiens, B., Helge, J.W., Richter, E.A. & Galbo, H. Interstitial glycerol concentrations in human skeletal muscle and adipose tissue during graded exercise, 367

© 2004 Scandinavian Physiological Society

Styf, J. *see* Zhang, Q.
Sun, Y., Fihn, B.-M., Jodal, M. & Sjövall, H. Inhibition of nitric oxide synthesis potentiates the colonic permeability increase triggered by luminal bile acids, 167
Syddall, H.E. *see* Holt, R.I.G.
Szentandrássy, N. *see* Fülöp, L.

Tan, H.L. *see* Verkerk, A.O.
Terkelsen, A.J., Andersen, O.K., Molgaard, H., Hansen, J. & Jensen, T.S. Mental stress inhibits pain perception and heart rate variability but not a nociceptive withdrawal reflex, 405
Tesch, P.A., Ekberg, A., Lindquist, D.M. & Trieschmann, J.T. Muscle hypertrophy following 5-week resistance training using a non-gravity-dependent exercise system, 89
Thompson, P.D. *see* Moyna, N.M.
Trieschmann, J.T. *see* Tesch, P.A.
Tsuchida, H. *see* Shibamoto, T.
Tupling, A.R. *see* Schertzer, J.D.

Vagianos, C.E. *see* Assimakopoulos, S.F.
Varró, A. *see* Fülöp, L.
Verkerk, A.O., Tan, H.L. & Ravesloot, J.H. Ca^{2+}-activated Cl^- current reduces transmural electrical heterogeneity within the rabbit left ventricle, 239

Wada, M. *see* Yamada, T.
Williams, J.H. Effects of fatigue on depolarization- and caffeine-induced contractures of skinned fibres, 265
Wittwer, M., Billeter, R., Hoppeler, H. & Flück, M. Regulatory gene expression in skeletal muscle of highly endurance-trained humans, 217
Wood, P.J. *see* Holt, R.I.G.
Wretenberg, P. *see* Al-Khalili, L.

Yamada, T., Inashima, S., Matsunaga, S., Nara, I., Kajihara, H. & Wada, M. Different time course of changes in sarcoplasmic reticulum and myosin isoforms in rat soleus muscle at early stage of hyperthyroidism, 79
Yamamoto, S.-I. *see* Kawashima, N.
Yano, H. *see* Kawashima, N.
Yaraş, N. *see* Yargiçoğlu, P.
Yargiçoğlu, P., Yaraş, N., Ağar, A., Gümüşlü, S., Abidin, İ. & Bilmen, S. Effects of *N*-nitro L-arginine methyl ester (L-NAME), a potent nitric oxide synthase inhibitor, on visual evoked potentials of rats exposed to different experimental stress models, 307
Yndestad, A. *see* Florholmen, G.
Yokoi, M. *see* Aoki, K.

Zhang, Q., Andersson, G., Lindberg, L.G. & Styf, J. Muscle blood flow in response to concentric muscular activity vs. passive venous compression, 57

© 2004 Scandinavian Physiological Society

Subject index

ABT-627, 319
action potential, 39
activation, 187
active hyperaemia, 231
adenosine, 301
adhesion, 143
adipose tissue, 367
adult height, 359
ageing, 99
Akt/protein kinase B, 395
ankle joint dorsiflexion and plantar flexion, 57
anterior pituitary cells, 347
arrhythmia, 13
arteriole, 231
α-subunit isoform, 49
attention to pain, 405

β_2-adrenergic agonist, 271
barbiturate, 29
bile acids, 167
bile duct ligation, 177
birthweight, 359
BK channels, 347

calcium, 265
calcium current, 39
calcium signalling, 335
Ca^{2+}-activated Cl^- current, 239
Ca^{2+}-ATPase, 195
capillary distensibility, 291
capillary endothelial cell, 291
Ca^{2+}-release, 195
Ca^{2+}-uptake, 195
cardiac output, 319
cardiac surgery, 5
cDNA filter arrays, 133
cell culture, 395
cell signalling, 143
cell volume, 329
circulation, 319
colonic mucosa, 167
concentric and eccentric muscle actions, 89
connective tissue, 125
coronary artery disease, 113
coronary stenosis, 21
cytokine, 133
cytoskeleton, 125

dispersion, 13
distraction from pain, 405
dopamine, 379
double occlusion pressure, 255

early afterdepolarization, 13
ECIS, 335
EDL, 209
elderly, 113
electrophysiology, 347
endothelial cell calcium, 231
endothelium, 113
endotoxaemia, 177
energetics, 187
energy metabolism, 133
enteric nervous system, 167
ERK mitogen-activated protein kinase, 395
excitability, 209
exercise, 71, 113, 195
exercise countermeasures, 89
exercise intensity, 367
exercise training, 157

fatigue, 209
flow cytometry, 143
flow heterogeneity, 21
flywheel technology, 89
football, 143
functional overload, 281

gain modulation, 99
gap junctions, 335
gene expression, 133, 217
GH_4 cells, 347
glycerol, 5, 367
growth hormone, 359
guinea-pig, 255

haemorrhage, 319
halothane, 29
heart, 157
heart failure, 113
heart rate variability, 405
heat shock protein 71, 72, 157, 387
hepatic circulation, 255
heterogeneity, 239
hindlimb suspension, 387
histamine receptor, 167
human experimental, 301
human myocytes, 39
humans, 405

iberiotoxin, 347
immunohistochemistry, 281
infancy, 359
inflammation, 125
insulin-like growth factor-I, 359
interferon-γ, 379
interstitial fluid pressure, 125
intestinal inflammation, 379
intestine, 177
iron scavanger, 21
ischaemia, 301
ischaemia-reperfusion injury, 21
isolated perfused liver, 255

knee extension torque, 249

© 2004 Scandinavian Physiological Society

lactate/pyruvate ratio, 5
L-DOPA, 379
length–tension relationship, 249
lipid peroxidation, 177, 307
lipolysis, 367
Losartan, 319

MDCK cells, 335
mechano-electrical feedback, 13
mental pressor test, 405
metabolic dilation, 231
microarray, 217
microdialysis, 5, 367
morphometric analysis, 291
muscle biopsy, 217, 367
muscle fibre, 217, 387
muscle fibre damage, 291
muscle function, 187
muscle plasticity, 271
muscle pump, 57
muscle strength and volume, 89
muscle-tendon unit, 249
M-waves, 209
myocardial infarction, 5
myocardial ischaemia, 5
MyoD, 281
myogenic regulatory factor, 271
myogenin, 281
myonuclei, 281
myosin heavy chain, 79
myosin light chain, 79

Na,K-ATPase, 49
Na^+,K^+-ATPase, 329
neonatal cardiomycoytes, 133
nitric oxide, 167, 307
N-nitro L-arginine methyl ester, 307
nociceptive withdrawal reflex, 405

obstructive jaundice, 177
oedema, 125
oscillations, 335
ouabain-insensitive, 49
over training, 143
oxidative capacity, 71
oxidative stress, 177
oxidative stress markers, 157
oxygen radicals, 21

p21, 281
pacing, 13
pain, 301
passive stiffness, 249
patch-clamp technique, 239
PCNA, 281
phalloidin, 125
photoplethysmography, 57
plantaris, 71
^{32}P-phosphorylation capacity, 49
prostaglandin E_1, 125
protein kinase C, 329
protein oxidation, 177

rat, 255
rat α_1, 49
recovery, 195
regional differences, 39
regulatory volume decrease, 329
renal excretion, 319
resistance vessels, 49
ryanodine receptor, 265

sarcoplasmic reticulum, 265
sarcoplasmic reticulum Ca^{2+}-ATPase, 79
satellite cell, 281
shock, 319
simulated haemorrhage, 29
sinusoidal pressure, 255
skeletal muscle, 231, 265, 367
soleus, 71, 209
spaceflight, 89
species difference, 255
spontaneously hypertensive rat, 29
Sprague–Dawley rat, 29
stochastic, 217
stress, 307
stretch, 13
stretch reflex, 99
superoxide dismutase, 157

theophylline, 301
thiol redox state, 177
thyroid hormone, 79
thyrotropin-releasing hormone, 347
tourniquet test, 301
training, 113
treadmill exercise, 387
type-I fibres, 209
type-II fibres, 209

urethane, 29

variability, 217
visual evoked potentials, 307
voltage clamp, 39
voluntary contraction, 99

Western blot, 49
Wistar–Kyoto rat, 29

xenotransplantation, 255

© 2004 Scandinavian Physiological Society